



## Coherus Announces Six-Year JUPITER-02 Follow-up Results Showing LOQTORZI® plus Chemotherapy Nearly Doubles Median Overall Survival in Nasopharyngeal Carcinoma

### -6 Years Long-term survival data reinforce potential benefit to patient survival in treating recurrent or metastatic nasopharyngeal carcinoma with LOQTORZI in combination with chemotherapy -

REDWOOD CITY, Calif., Dec. 08, 2025 (GLOBE NEWSWIRE) -- Coherus Oncology, Inc. (NASDAQ: CHRS) today announced compelling six-year overall survival (OS) follow-up results from the Phase 3 JUPITER-02 trial evaluating LOQTORZI® (toripalimab-tpzi) plus chemotherapy in recurrent or metastatic nasopharyngeal carcinoma (RM-NPC). The findings reveal a striking and durable survival advantage that underscores the urgent clinical need to incorporate LOQTORZI with chemotherapy as first-line treatment.

In this exploratory post-hoc analysis, patients receiving LOQTORZI plus gemcitabine and cisplatin achieved a median OS of 64.8 months, nearly double that of chemotherapy alone (33.7 months), representing a 31-month improvement and an observed 38% reduction in risk of death (HR 0.62; 95% CI, 0.45–0.85). These results, presented at ESMO Asia 2025, signal a step change in cancer patient survival, reinforcing LOQTORZI's role in transforming outcomes for people living with RM-NPC.

JUPITER-02 is a randomized, double-blind, placebo-controlled Phase 3 study evaluating LOQTORZI with chemotherapy in first-line RM-NPC, and this long-term follow-up provides additional context for the previously reported survival outcomes.

#### A Meaningful Shift for Patients Who Need It Most

RM-NPC is an aggressive cancer, and long-term survival with standard chemotherapy can be limited for many patients. The multi-year survival observed in the LOQTORZI arm suggests a potential for meaningful clinical benefit, which may translate into longer survival for patients who typically face a challenging prognosis.

"The new 6-year overall survival follow up data gives us even greater confidence to use toripalimab in patients with NPC that is recurrent or metastatic," said Victoria Villaflor, MD, Professor and Director, Head and Neck Oncology Program, Division of Hematology-Oncology, Department of Medicine, UC Irvine School of Medicine.

For many patients, the difference between 33 months and nearly 65 months represents the possibility of more time with family and more milestones. This meaningful extension highlights why oncologists may consider adding LOQTORZI to chemotherapy upfront, as delaying or omitting a therapy associated with improved survival outcomes could reduce a patient's opportunity to achieve longer-term benefit.

#### A Standard of Care Reinforced by Long-Term Evidence

"These data suggest a significant long-term overall survival benefit for patients living with RM-NPC," said Rosh Dias, MD, Chief Medical Officer, Coherus Oncology. "With these long-term data, LOQTORZI, in combination with chemotherapy, reinforces the data supporting this regimen as the standard of care for patients living with RM-NPC."

Coherus Oncology is advancing a pipeline built on deep scientific expertise and strategic collaborations designed to deliver first- and best-in-class therapies. LOQTORZI, Coherus' next-generation PD-1 inhibitor, is an important part of this vision, with data indicating its potential to enhance survival outcomes when used with chemotherapy.

#### ESMO Asia 2025 Presentation Details

**Abstract # 1279:** [Long Term Overall Survival Follow-up of Toripalimab versus Placebo in Combination with Gemcitabine and Cisplatin as First-line Treatment for Recurrent or Metastatic Nasopharyngeal Carcinoma](#)

- Date: Friday, December 5, 2025, 5:00 p.m. – 6:30 p.m. PST

#### INDICATIONS AND IMPORTANT SAFETY INFORMATION

##### INDICATIONS

LOQTORZI® (toripalimab-tpzi) is indicated:

- In combination with cisplatin and gemcitabine, for the first-line treatment of adults with metastatic or with recurrent, locally advanced nasopharyngeal carcinoma (NPC).
- As a single agent, for the treatment of adults with recurrent unresectable or metastatic NPC with disease progression on or after a platinum-containing chemotherapy.

##### IMPORTANT SAFETY INFORMATION

###### Severe and Fatal Immune-Mediated Adverse Reactions

Immune-mediated adverse reactions listed herein may not include all possible severe and fatal immune-mediated adverse reactions. Immune-mediated adverse reactions, which can be severe or fatal, occur in any organ system or tissue, affect more than one body system simultaneously, and occur at any time after starting PD-1/PD-L1 blocking antibody. While immune-mediated adverse reactions usually manifest during treatment, they can also manifest after discontinuation of PD-1/PD-L1 blocking antibodies.

- Monitor for early identification and management. Evaluate liver enzymes, creatinine, and thyroid function at baseline and periodically during treatment. In cases of suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation

as appropriate.

- Withhold or permanently discontinue LOQTORZI based on severity and type of reaction (see Dosage and Administration in Prescribing Information). In general, if LOQTORZI requires interruption or discontinuation, administer systemic corticosteroid therapy (1 to 2 mg/kg/day prednisone or equivalent) until improvement to Grade 1 or less. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reactions are not controlled with corticosteroid therapy.

#### Immune-Mediated Pneumonitis

LOQTORZI can cause immune-mediated pneumonitis.

- In patients receiving LOQTORZI in combination with cisplatin and gemcitabine, immune-mediated pneumonitis occurred in 2.1% (3/146) of patients, including Grade 2 (1.4%) adverse reactions. Pneumonitis resolved in 67% (2/3) of these patients.
- In patients receiving LOQTORZI monotherapy, immune-mediated pneumonitis occurred in 2.6% (22/851) of patients, including fatal (0.2%), Grade 3 (0.7%), and Grade 2 (1.1%) adverse reactions. Systemic corticosteroids were required in 82% (18/22) of patients with pneumonitis. Pneumonitis led to permanent discontinuation of LOQTORZI in 1.2% (10/851) of patients. Pneumonitis resolved in 23% (5/22) of these patients.

#### Immune-Mediated Colitis

LOQTORZI can cause immune-mediated colitis, which may present with diarrhea. Cytomegalovirus (CMV) infection/reactivation has been reported in patients with corticosteroid-refractory immune-mediated colitis. In cases of corticosteroid-refractory colitis, consider repeating infectious workup to exclude alternative etiologies. In patients receiving LOQTORZI monotherapy, immune-mediated colitis occurred in 0.4% (3/851) of patients, including Grade 3 (0.2%) and Grade 2 (0.1%) adverse reactions. Colitis resolved in all 3 patients.

#### Hepatotoxicity and Immune-Mediated Hepatitis

LOQTORZI can cause immune-mediated hepatitis.

- In patients receiving LOQTORZI in combination with cisplatin and gemcitabine, immune-mediated hepatitis occurred in 0.7% (1/146) of patients, which was a Grade 3 (0.7%) adverse reaction. The patient with immune-mediated hepatitis required systemic corticosteroids.
- In patients receiving LOQTORZI monotherapy, immune-mediated hepatitis occurred in 3.3% (28/851) of patients, including Grade 4 (0.8%), Grade 3 (2.1%), and Grade 2 (0.4%) adverse reactions. Hepatitis led to permanent discontinuation of LOQTORZI in 1.1% of patients and withholding of LOQTORZI in 0.8% of patients. Hepatitis resolved in 54% (15/28) of these patients.

#### Immune-Mediated Endocrinopathies

##### *Adrenal Insufficiency*

LOQTORZI can cause primary or secondary adrenal insufficiency. For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment, including hormone replacement as clinically indicated. Withhold or permanently discontinue LOQTORZI depending on severity. In patients receiving LOQTORZI monotherapy, adrenal insufficiency occurred in 0.5% (4/851) of patients, including Grade 2 (0.4%) and Grade 1 (0.1%) adverse reactions. Systemic corticosteroids were required in 75% (3/4) of the patients with adrenal insufficiency. Adrenal insufficiency led to withholding of LOQTORZI in 0.1% (1/851) of patients. In the one patient in whom LOQTORZI was withheld, LOQTORZI was reinitiated after symptom improvement.

##### *Hypophysitis*

LOQTORZI can cause immune-mediated hypophysitis. Hypophysitis can present with acute symptoms associated with mass effects such as headache, photophobia, or visual field defects. Hypophysitis can cause hypopituitarism. Initiate hormone replacement as indicated. Withhold or permanently discontinue LOQTORZI depending on severity. In patients receiving LOQTORZI monotherapy, hypophysitis occurred in 0.4% (3/851) of patients receiving LOQTORZI, including Grade 3 (0.2%) and Grade 2 (0.1%) adverse reactions. All three patients received systemic corticosteroids. Hypophysitis led to permanent discontinuation of LOQTORZI in 0.1% (1/851) of patients and withholding of LOQTORZI in 0.1% (1/851) of patients. The one patient in whom LOQTORZI was withheld reinitiated LOQTORZI.

##### *Thyroid Disorders*

LOQTORZI can cause immune-mediated thyroid disorders. Thyroiditis can present with or without endocrinopathy. Hypothyroidism can follow hyperthyroidism. Initiate hormone replacement for hypothyroidism or institute medical management of hyperthyroidism as clinically indicated. Withhold or permanently discontinue LOQTORZI depending on severity.

- In patients receiving LOQTORZI in combination with cisplatin and gemcitabine, thyroiditis occurred in 2.1% (3/146) of patients receiving LOQTORZI, including Grade 2 (1.4%). Three patients required thyroid hormone replacement therapy. Thyroiditis resolved in one of the 3 patients. Hyperthyroidism occurred in 1.4% (2/146) of patients receiving LOQTORZI in combination with cisplatin and gemcitabine. Hyperthyroidism resolved in these 2 patients. Hypothyroidism occurred in 30% (44/146) of patients receiving LOQTORZI in combination with cisplatin and gemcitabine, including Grade 2 (24%) and Grade 1 (6%). Eighty percent of the 44 patients required thyroid hormone replacement therapy. LOQTORZI was withheld in 2.1% (3/146) of the patients. Of the 3 patients in whom LOQTORZI was withheld, 2 patients reinitiated LOQTORZI.
- In patients receiving LOQTORZI monotherapy, thyroiditis occurred in 0.6% (5/851) patients receiving LOQTORZI, including Grade 2 (0.1%). Two of these 5 patients received systemic corticosteroids and 2 required thyroid hormone replacement therapy. Thyroiditis resolved in 2 of the 5 patients. Hyperthyroidism occurred in 7% (55/851) of patients receiving LOQTORZI, including Grade 2 (1.9%). Hyperthyroidism resolved in 85% (47/55) of the patients. Hypothyroidism occurred in 15% (128/851) of patients receiving LOQTORZI, including Grade 2 (8%). Sixty three percent of the 128 patients required thyroid hormone replacement therapy. LOQTORZI was withheld in 0.5% of patients. Of the 4 patients in whom LOQTORZI was withheld, 3 patients reinitiated LOQTORZI.

#### *Type 1 Diabetes Mellitus, which can present with Diabetic Ketoacidosis*

Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold or permanently discontinue LOQTORZI depending on severity. In patients receiving LOQTORZI monotherapy, diabetes mellitus occurred in 0.9% (8/851) of patients receiving LOQTORZI, including Grade 4 (0.1%), Grade 3 (0.7%), and Grade 2 (0.1%). Diabetes mellitus led to permanent discontinuation in 0.4% of patients. Six of the 8 (75%) patients with diabetes mellitus required long-term insulin therapy.

#### Immune-Mediated Nephritis with Renal Dysfunction

LOQTORZI can cause immune-mediated nephritis.

- In patients receiving LOQTORZI in combination with cisplatin and gemcitabine, immune-mediated nephritis occurred in 0.7% (1/146) of patients receiving LOQTORZI. The one patient with immune-mediated nephritis (Grade 4) required systemic corticosteroids and nephritis led to discontinuation of LOQTORZI. Nephritis resolved in this patient.
- In patients receiving LOQTORZI monotherapy, immune-mediated nephritis occurred in 0.5% (4/851) of patients, including Grade 3 (0.5%) adverse reactions. Nephritis resolved in 75% (3/4) of these patients.

#### Immune-Mediated Dermatologic Adverse Reactions

LOQTORZI can cause immune-mediated rash or dermatitis. Exfoliative dermatitis, including Stevens-Johnson Syndrome (SJS), drug rash with eosinophilia and systemic symptoms (DRESS), and toxic epidermal necrolysis (TEN), has occurred with PD-1/PD-L1 blocking antibodies. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-exfoliative rashes. Withhold or permanently discontinue LOQTORZI depending on severity.

- In patients receiving LOQTORZI in combination with cisplatin and gemcitabine, immune-mediated dermatologic adverse reactions occurred in 8% (12/146) of patients, including Grade 3 (3.4%) and Grade 2 (1.4%) adverse reactions. Systemic corticosteroids were required in 25% (3/12) of the patients with immune-mediated dermatologic adverse reactions. Immune-mediated dermatologic adverse reactions led to permanent discontinuation of LOQTORZI in 2.1% (3) of patients. Immune-mediated dermatologic adverse reactions resolved in 92% (11/12) of these patients.
- In patients receiving LOQTORZI monotherapy, immune-mediated dermatologic adverse reactions occurred in 4% (34/851) of patients, including Grade 3 (0.4%) and Grade 2 (1.4%) adverse reactions. Immune-mediated dermatologic adverse reactions led to withholding of LOQTORZI in 0.4% (3) of the patients. Systemic corticosteroids were required in 12% (4/34) of the patients with immune-mediated dermatologic adverse reactions. Immune-mediated dermatologic adverse reactions resolved in 71% (24/34) of these patients.

#### Other Immune-Mediated Adverse Reactions

The following clinically significant immune-mediated adverse reactions occurred at an incidence of <1% (unless otherwise noted) in patients who received LOQTORZI or were reported with the use of other PD-1/PD-L1 blocking antibodies. Severe or fatal cases have been reported for some of these adverse reactions.

- **Cardiac/Vascular:** Myocarditis, pericarditis, vasculitis, pericardial effusion
- **Nervous System:** Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis (including exacerbation), Guillain-Barré syndrome, nerve paresis, autoimmune neuropathy
- **Ocular:** Uveitis, iritis and other ocular inflammatory toxicities can occur. Some cases can be associated with retinal detachment. Various grades of visual impairment, including blindness, can occur. If uveitis occurs in combination with other immune-mediated adverse reactions, consider a Vogt-Koyanagi-Harada-like syndrome, as this may require treatment with systemic steroids to reduce the risk of permanent vision loss.
- **Gastrointestinal:** Pancreatitis, to include increases in serum amylase and lipase levels, gastritis, duodenitis
- **Musculoskeletal and Connective Tissue:** Myositis/polymyositis, rhabdomyolysis (and associated sequelae, including renal failure), arthritis, polymyalgia rheumatica, dermatomyositis
- **Endocrine:** Hypoparathyroidism
- **Hematologic/Immune:** Hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis, systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenic purpura, solid organ transplant rejection, other transplant (including corneal graft) rejection

#### Infusion-Related Reactions

LOQTORZI can cause severe or life-threatening infusion-related reactions including hypersensitivity and anaphylaxis.

- In patients receiving LOQTORZI in combination with cisplatin and gemcitabine, infusion-related reactions have been reported in 4.1% of patients, including Grade 2 (0.7%) reactions.
- In patients receiving LOQTORZI monotherapy, infusion-related reactions occurred in 2% of 851 patients, including Grade 3 (0.1%) and Grade 2 (0.6%). LOQTORZI was withheld for one Grade 3 infusion related reaction. Monitor patients for signs and symptoms of infusion-related reactions including rigors, chills, wheezing, pruritus, flushing, rash, hypotension, hypoxemia, and fever. Interrupt or slow the rate of infusion for mild (Grade 1) or moderate (Grade 2) infusion-related reactions. For severe (Grade 3) or life-threatening (Grade 4) infusion-related reactions, stop infusion and permanently discontinue LOQTORZI.

#### Complications of Allogeneic Hematopoietic Stem Cell Transplant (HSCT)

Fatal and other serious complications can occur in patients who receive allogeneic hematopoietic stem cell transplantation (HSCT) before or after being treated with a PD-1/PD-L1 blocking antibody. Transplant-related complications include hyperacute graft-versus-host-disease (GVHD), acute GVHD, chronic GVHD, hepatic veno-occlusive disease (VOD) after reduced intensity conditioning, and steroid requiring febrile syndrome (without an identified infectious cause). These complications may occur despite intervening therapy between PD-1/PD-L1 blockade and allogeneic HSCT. Follow patients closely for evidence of transplant-related complications and intervene promptly. Consider the benefit versus risks of treatment with a

PD-1/PD-L1 blocking antibody prior to or after an allogeneic HSCT.

### Embryo-Fetal Toxicity

LOQTORZI can cause fetal harm when administered to a pregnant woman. Animal studies have demonstrated that inhibition of the PD-1/PD-L1 pathway can lead to increased risk of immune-mediated rejection of the developing fetus resulting in fetal death. Advise women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with LOQTORZI and for 4 months after the last dose.

### Lactation

There are no data on the presence of toripalimab-tpzi in human milk; its effects on the breastfed child, or on milk production. Maternal IgG is known to be present in human milk. The effects of local gastrointestinal exposure and limited systemic exposure in the breastfed child to toripalimab-tpzi are unknown. Because of the potential for serious adverse reactions in breastfed children, advise lactating women not to breastfeed during treatment with LOQTORZI and for 4 months after the last dose.

### Serious Adverse Reactions

- In JUPITER-02, when LOQTORZI was administered in combination with cisplatin and gemcitabine for the first-line treatment of recurrent, locally advanced or metastatic nasopharyngeal carcinoma, serious adverse reactions occurred in 43% of patients. Serious adverse drug reactions in  $\geq 2\%$  were thrombocytopenia (14%), neutrophil count decreased (10%), pneumonia (10%), anemia (9%), abnormal hepatic function (2.7%), and rash (2.1%). There were three fatal adverse reactions (2.1%): one due to epistaxis; one due to intracranial hemorrhage associated with immune-related thrombocytopenia and coagulopathy; and one due to pneumonia. Permanent discontinuation of LOQTORZI, due to an adverse reaction occurred in 12% of patients. Adverse reactions resulting in permanent discontinuation of LOQTORZI in  $\geq 1\%$  were pneumonia (2.1%), pulmonary tuberculosis (1.4%), rash (1.4%), and vomiting (1.4%). The most common Grade 3 to 4 laboratory abnormalities ( $\geq 2\%$ ) were decreased neutrophils (58%), decreased lymphocytes (57%), decreased hemoglobin (50%), decreased platelets (33%), decreased potassium (10%), decreased sodium (9%), increased alanine aminotransferase (6%), increased or decreased magnesium (4.2% each), decreased calcium (3.5%), increased aspartate aminotransferase (2.7%), and increased bilirubin (2.1%).
- In POLARIS-02, when LOQTORZI was administered as a single agent to patients with previously treated, unresectable or metastatic nasopharyngeal carcinoma, serious adverse reactions occurred in 24% of patients. Serious adverse drug reactions in  $\geq 2\%$  were pneumonia (4.7%), abnormal hepatic function (2.6%), and hyperbilirubinemia (2.1%). Fatal adverse reactions occurred in 3.7% of patients who received LOQTORZI, including death not otherwise specified (1.6%), tumor hemorrhage (0.5%), hepatic failure and thrombocytopenia (0.5%), hyponatremia (0.5%), and sudden death (0.5%). Permanent discontinuation of LOQTORZI due to an adverse reaction occurred in 9% of patients. Adverse reactions resulting in permanent discontinuation of LOQTORZI in  $\geq 1\%$  included pneumonia (1.1%), abnormal hepatic function (1.1%), and hyperbilirubinemia (1.1%). The most common Grade 3 or 4 laboratory abnormalities ( $\geq 2\%$ ), were decreased sodium (11%), decreased lymphocytes (9%), decreased hemoglobin (6%), increased aspartate aminotransferase (3.8%), decreased phosphate (3.2%), and increased alkaline phosphatase (2.2%).

### Common Adverse Reactions

- In JUPITER-02, the most common adverse reactions ( $\geq 20\%$ ) were nausea (71%), vomiting (68%), decreased appetite (55%), constipation (39%), hypothyroidism (38%), rash (36%), pyrexia (32%), diarrhea (31%), peripheral neuropathy (30%), cough (26%), musculoskeletal pain (25%), upper respiratory infection (23%), insomnia (23%), dizziness (21%), and malaise (21%).
- In POLARIS-02, in patients with previously treated, unresectable or metastatic nasopharyngeal carcinoma, the most common ( $\geq 20\%$ ) adverse reactions were hypothyroidism (27%), fatigue (22%), and cough (20%).

LOQTORZI® Injection: 240 mg/6 mL (40 mg/mL) solution in a single-dose vial

Please see [prescribing information](#) for LOQTORZI.

### About Coherus Oncology

Coherus Oncology is a fully integrated commercial-stage innovative oncology company with an approved next-generation PD-1 inhibitor, LOQTORZI® (toripalimab-tpzi), and a pipeline that includes two mid-stage clinical candidates targeting liver, lung, head & neck, colorectal and other cancers. The Company's strategy is to grow sales of LOQTORZI in NPC and advance the development of new indications for LOQTORZI in combination with both its pipeline candidates as well as through its partners, driving sales multiples and synergies from proprietary combinations.

Coherus' innovative oncology pipeline includes multiple antibody immunotherapy candidates focused on enhancing the innate and adaptive immune responses to enable a robust antitumor response and enhance outcomes for patients with cancer. CHS-114 is a highly selective cytolytic anti-CCR8 antibody currently in Phase 1b/2a studies in patients with advanced solid tumors, including head and neck squamous cell carcinoma, colorectal cancer, gastric cancer, and esophageal cancer. Casdozokitug is a novel IL-27 antagonistic antibody currently being evaluated in a Phase 2 study in patients with first-line hepatocellular carcinoma.

For more information about LOQTORZI, including the U.S. Prescribing Information and important safety information, please visit [www.loqtorzi.com](http://www.loqtorzi.com).

### Forward-Looking Statements

The statements in this press release include express or implied forward-looking statements within the meaning of the "safe harbor" provisions of the Private Securities Litigation Reform Act of 1995, Section 27A of the Securities Act 1933, as amended, and Section 21E of the Securities Exchange Act of 1934, as amended about Coherus that involve risks and uncertainties relating to future events and the future performance of Coherus. Forward-looking statements relate to expectations, beliefs, projections, future plans and strategies, anticipated events or trends and similar expressions concerning matters that are not historical facts. Words such as "will," "could," "would," "should," "expect," "plan," "anticipate," "intend," "believe," "estimate," "predict," "project," "potential," "continue," "future," "opportunity," "likely," "target," variations of such words, and similar expressions or

negatives of these words are intended to identify such forward-looking statements, although not all forward-looking statements contain these identifying words. You can also identify forward-looking statements by discussions of strategy, plans or intentions.

Examples of such forward-looking statements include, but are not limited to, express or implied statements regarding: the ability of Coherus' pipeline to enhance outcomes for cancer patients; expectations about future synergies; projections about growth in sales; expectations for future enrollment in clinical trials; projections about the expansion of indications for LOQTORZI; and the assumptions underlying or relating to such statements.

These forward-looking statements are based on Coherus' current plans, estimates and projections. Such forward-looking statements involve substantial risks and uncertainties that could cause actual results to differ materially from those contemplated in any forward-looking statements. Such risks and uncertainties include, without limitation: uncertainties about the potential impact of unforeseen liabilities, future capital expenditures, revenues, costs, expenses, earnings, economic performance, indebtedness, financial condition and losses on Coherus' prospects, business and operations in the future; risks and uncertainties in executing collaboration agreements and other joint ventures; risks and uncertainties of conducting clinical trials; the risks of Coherus' dependence on an ability to raise funds, which may not be available on acceptable terms or at all; and risks and uncertainties of any litigation, regulatory actions and other legal proceedings.

All forward-looking statements contained in this press release speak only as of the date of this press release. Coherus undertakes no obligation to update or revise any forward-looking statements. For a further discussion of these and other factors that could cause Coherus' future results to differ materially from any forward-looking statements see the section entitled "Risk Factors" in Coherus' Quarterly Report on Form 10-Q for the period ended September 30, 2025, filed with the Securities and Exchange Commission (SEC) on November 6, 2025, as updated by Coherus' subsequent reports filed with the SEC.

LOQTORZI® is a registered trademark of Coherus Oncology, Inc.  
©2025 Coherus Oncology, Inc. All rights reserved.

**Coherus Oncology Contact Information:**

For Investors:  
Carrie Graham  
VP, Investor Relations & Advocacy  
[IR@coherus.com](mailto:IR@coherus.com)

US-LOQ-0283.1 12/2025



 [Primary Logo](#)

Source: Coherus Oncology, Inc.